

## R-H. Hadron physics

Wigner research group

**Ferenc Siklér**, László Boldizsár, Zoltán Fodor<sup>A</sup>, Endre Futó<sup>A</sup>, Sándor Hegyi, Gábor Jancsó<sup>A</sup>, József Kecskeméti<sup>A</sup>, Krisztián Krajczár, Andrew John Lowe, Gabriella Pála<sup>E</sup>, Sona Pochybová<sup>#</sup>, Zoltán Seres<sup>A</sup>, Anna Julia Zsigmond<sup>#</sup>



**Quarks and gluons.** — Particle physics is our attempt to understand the basic constituents of our world. What is it made of? What are the interactions between the building blocks of matter? Symmetries and gauge theories provide a coherent framework for the electromagnetic, weak, and strong interactions. The last of these, the strong force, acts between quarks and gluons and is described by the theory of quantum chromodynamics (QCD). In most circumstances, it is difficult to perform accurate calculations with QCD because the theory is strongly coupled and consequently has a nonperturbative nature. Results from the Relativistic Heavy Ion Collider (RHIC) at Brookhaven National Laboratory, later reinforced by those from the Large Hadron Collider (LHC) at CERN, showed unexpected phenomena: suppression of hadrons with high transverse momentum ( $p_T$ ), and weakening of back-to-back jet correlations. These results indicated that quark matter does not behave as a quasi-ideal state of free quarks and gluons, but as an almost perfect dense fluid.

Our research group studies collisions of nucleons and nuclei, performs basic and advanced measurements, and tests theoretical ideas. We participate in several complementary experiments (ALICE and CMS), both in data-taking and physics analysis. In the past year our research group concentrated mostly on the analysis of pPb data recorded at LHC at  $\sqrt{s_{NN}} = 5.02$  TeV energy per nucleon pair. The large amount of collected data allowed us to perform the studies proposed at the beginning of the year.

**Size and shape of the created system in pPb collisions.** — Measurements of the correlation between hadrons emitted in high-energy collisions of nucleons and nuclei can be used to study the spatial extent and shape of the created system. The characteristic radii and the homogeneity lengths of the particle-emitting source can be extracted with reasonable precision. We have studied the characteristics of the one-, two-, and three-dimensional two-particle correlation functions in various center-of-mass energy pp, pPb, and peripheral PbPb collisions as a function of the transverse pair momentum  $k_T$  and of the charged-particle multiplicity  $N_{\text{tracks}}$  of the event. Charged pions and kaons at low  $p_T$  and in laboratory pseudorapidity  $|\eta| < 1$  were identified via their energy loss in the silicon tracker. The correlation functions were corrected for the Coulomb interaction between particles. The contributions from other, correlated particle emissions (mini-jets, multi-body resonance decays) were also subtracted. The obtained distributions could be fit by an exponential parametrization in the relative momentum of the particle pair, both in one- and multi-dimensions.


---

<sup>A</sup> Associate fellow

<sup>E</sup> Professor Emeritus

<sup>#</sup> Ph.D. student

The extracted exponential radii for pions increase with increasing  $N_{\text{tracks}}$  for all systems and center-of-mass energies studied, for one, two, and three dimensions alike. Their values are in the range 1–5 fm, reaching their highest values for very high multiplicity pPb and for similar multiplicity PbPb collisions. The  $N_{\text{tracks}}$  dependence of longitudinal ( $R_l$ ) and transverse radii ( $R_t$ ) is similar for pp and pPb in all  $k_T$  bins, and that similarity also applies to peripheral PbPb if  $k_T > 0.4$  GeV/c. In general there is an ordering for the radii:  $R_l > R_t$ , thus the pp and pPb source is elongated in the beam direction. In the case of peripheral PbPb the source is quite symmetric, and shows a slightly different  $N_{\text{tracks}}$  dependence. The most visible divergence between pp, pPb, and PbPb is seen in the so-called “out” radius ( $R_o$ ) that could indicate the differing lifetime of the created systems in those collisions.




**Figure 1. Left:** The like-sign correlation function of pions (red triangles) corrected for Coulomb interaction and cluster contribution (mini-jets and multi-body resonance decays) as a function of the combined momentum, in a selected  $N_{rec}$  bins for all  $k_T$ . The solid curve indicates a fit with the exponential Bose-Einstein parametrization. **Right:** The longitudinal radius  $R_l$  as a function of  $N_{\text{tracks}}$  scaled to  $k_T = 0.45$  GeV/c with help of a parametrization.

The kaon radii also show some increase with  $N_{\text{tracks}}$ , although its magnitude is smaller than that for pions. Longer-lived resonances and rescattering may play a role here. The pion radii decrease with increasing  $k_T$ . The dependence of the radii on the multiplicity and  $k_T$  factorizes and in some cases appears to be less sensitive to the type of the colliding system and center-of-mass energy. The similarities observed in the  $N_{\text{tracks}}$  dependence may point to a common critical hadron density in pp, pPb, and peripheral PbPb collisions, since the present correlation technique measures the characteristic size of the system near the time of the last interactions.

**Spectra of high  $p_T$  charged hadrons in pPb collisions.** — We have measured the charged-particle spectra in pPb collisions in the transverse momentum range of  $0.4 < p_T < 120$  GeV/c for center-of-mass pseudorapidities up to  $|\eta_{CM}| = 1.8$ . The forward-backward yield asymmetry has been measured as a function of  $p_T$  for three bins in  $\eta_{CM}$ . At  $p_T < 10$  GeV/c, the charged-particle production is enhanced in the direction of the Pb beam, in qualitative agreement with nuclear shadowing expectations. The nuclear modification factor at mid-

rapidity, relative to a reference spectrum interpolated from pp measurements at lower and higher collision energies, rises above unity at high  $p_T$  reaching an  $R_{pPb}$  value of 1.3–1.4 at  $p_T \geq 40$  GeV/c. The observed enhancement is larger than expected from next-to-leading order (NLO) perturbative QCD predictions that include anti-shadowing effects in the nuclear parton distribution functions (nPDFs) in this kinematic range.




**Figure 2. Left:** Charged-particle forward-backward yield asymmetry as a function of  $p_T$  for three  $|\eta_{CM}|$  intervals. The asymmetry is computed as the charged-particle yields in the direction of the Pb beam divided by those of the proton beam. **Right:** Charged-particle nuclear modification factors measured by CMS in  $|\eta_{CM}| < 1$  (filled circles), and by ALICE in  $|\eta_{CM}| < 0.3$  (open squares), are compared to a theoretical prediction.

The fact that the nuclear modification factor is below unity for  $p_T < 2$  GeV/c is anticipated since particle production in this region is dominated by softer scattering processes that are not expected to scale with the nuclear thickness function. In the intermediate  $p_T$  range (2–5 GeV/c), no significant deviation from unity is found in the  $R_{pPb}$  ratio. There are several prior measurements that suggest an interplay of multiple effects in this  $p_T$  region. At lower collision energies, an enhancement (“Cronin effect”) has been observed that is larger for baryons than for mesons, and is stronger in the more central collisions. This enhancement has been attributed to a combination of initial-state multiple scattering effects, causing momentum broadening, and hadronization through parton recombination (a final-state effect) invoked to accommodate baryon/meson differences. Recent results from pPb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV and from dAu collisions at  $\sqrt{s_{NN}} = 200$  GeV suggest that collective effects may also play a role in the intermediate- $p_T$  region. Most theoretical models do not predict a Cronin enhancement in this  $p_T$  range at LHC energies as the effect of initial-state multiple scattering is compensated by nPDF shadowing.

The observed rise of the nuclear modification factor up to  $R_{pPb} \approx 1.3\text{--}1.4$  at high  $p_T$ , albeit with large uncertainty, is much stronger than expected theoretically. None of the available theoretical models predict enhancements beyond  $R_{pPb} \approx 1.1$  at high  $p_T$ . In particular, although the  $p_T$  range corresponds to parton momentum fractions  $0.02 \leq x \leq 0.2$ , which coincides with the region where parton anti-shadowing effects are expected, none of the nPDFs obtained from global fits to nuclear data predict enhancements beyond 10% at the large virtualities of relevance here. We also show the measurement of the ALICE Collaboration, which is performed in a narrower pseudorapidity range than the CMS one, and uses a different method to obtain the pp reference spectrum based on ALICE pp data measured at  $\sqrt{s} = 7$  TeV. The difference in the CMS and ALICE  $R_{pPb}$  results stems primarily from differences in the charged-hadron spectra measured in pp collisions at  $\sqrt{s} = 7$  TeV.

Future direct measurement of the spectra of jets and charged particles in pp collisions at a center-of-mass energy of 5.02 TeV is necessary to better constrain the fragmentation functions and also to reduce the dominant systematic uncertainties in the charged-particle nuclear modification factor.

**Spectra of high mass bosons in pPb collisions.** — We have measured the Z boson production cross section in the muon decay channel in pPb collisions. The results are presented in the center-of-mass frame with positive rapidity values corresponding to the proton fragmentation region. The Z boson candidates are selected as an opposite-charge muon pair in the 60–120 GeV/c<sup>2</sup> mass range where both muons have  $p_T > 20$  GeV/c and are within the  $|\eta_{lab}| < 2.4$  muon detector coverage. The measured inclusive Z boson production cross section in pPb collisions for the range  $-2.5 < y_{CM} < 1.5$  is  $\sigma_{pPb}(Z \rightarrow \mu\mu) = 94.1 \pm 2.1$  (stat.)  $\pm 2.4$  (syst.)  $\pm 3.3$  (lumi.) nb using the calibrated integrated luminosity. For the same restricted rapidity range, the POWHEG simulation predicts  $94.0 \pm 4.7$  nb after multiplying by the number of nucleons in the Pb nucleus ( $A = 208$ ), which corresponds to the hypothesis of binary collision scaling in pPb.



**Figure 3. Left:** Invariant mass of selected muon pairs from pPb data compared to a simulation that was normalized to the number of events in data in the signal mass regions. **Right:** Differential cross section of Z bosons in pPb collisions as a function of rapidity compared to predictions from POWHEG+PYTHIA generator with CT10NLO PDF set, from PYTHIA generator with Z2 underlying event tune. All theory predictions are scaled by  $A = 208$ .

The differential cross section as a function of Z boson rapidity is consistent with the theory predictions. The forward-backward ratio, defined as  $d\sigma(+y)/d\sigma(-y)$ , is expected to be more sensitive to nuclear effects, because normalization uncertainties cancel both in theory and in experiment. Due to the large statistical uncertainties, this measurement is unable to distinguish between different nPDF sets but it can constrain their uncertainties by adding new data points to the global fits in a previously unexplored region of the  $Q^2$ -x phase space. The differential cross section as a function of Z boson transverse momentum has been measured and apart from very low transverse momenta it is in good agreement with the predictions from PYTHIA. The results of the presented measurement provide new data points in a previously unexplored region of phase space for constraining nuclear PDF fits.

Thus, Z boson production is unmodified by the hot and dense QCD medium produced in heavy ion collisions, and its yield scales with the number of binary nucleon-nucleon collisions. The nuclear modification factor does not exhibit large deviations from unity showing that nuclear effects are small with respect to the uncertainties of the pPb measurements. The results were compared to NLO theory predictions with and without nuclear modification, that show hints of nuclear effects but more luminosity is needed to distinguish between different nPDF sets. These measurements set constraints for the global fits of nPDFs in a previously unexplored region of phase space.

**Quark and gluon jets.** — We have studied the proton-to-pion ratio in jets produced in simulated proton-proton collisions at  $\sqrt{s} = 7$  TeV using the PYTHIA 6.4 Monte Carlo (MC) event generator. We compared the  $p/\pi$  ratio in the selected quark-like and gluon-like jets to a reference sample of tagged quark- and gluon-jets. The contamination in the selected jets significantly influences the observed ratios. Thus, despite the different fragmentation of jets originating from quarks or gluons, we see no difference in the proton-to-pion ratio inside these jets, within the used MC model. To see whether this statement holds, we suggest proceeding with similar study using experimental data.

## Grants

OTKA K 81614: New analysis methods and tests of quantum chromodynamics at the LHC (F. Siklér, 2010-2014)

OTKA K 109703: Consortial main: Hungary in the CMS experiment of the Large Hadron Collider (F. Siklér, 2013-2016)

Swiss National Science Foundation, SCOPES 152601: Preparation for and exploitation of the CMS data taking at the next LHC run (G. Dissertori ETHZ, 2014-2017)

EC FP7 C 262025: Advanced European Infrastructures for Detectors and Accelerators [AIDA] (F. Siklér, 2011-2014)

“Wigner research group” support (F. Siklér, 2014)

## Publications

### Articles

1. Münzer R et al. incl. [Fodor Z](#), [Kecskemeti J](#), [Seres Z](#) (67 authors): SiAvio: A trigger for  $\Lambda$ -hyperons. **NUCL INSTRUM METH A**, 745: pp. 38-49. (2014)

See also: R-I.3,

### CMS collaboration

Due to the vast number of publications of the large collaborations in which the research group participated in 2014, here we list only a short selection of appearances in journals with the highest impact factor.

1. Chatrchyan S et al. incl. [Bencze Gy](#), [Hajdu Cs](#), [Hidas P](#), [Horvath D](#), [Sikler F](#), [Veszpremi V](#), [Vesztergombi Gy](#), [Zsigmond AJ](#) [2200 authors]: Evidence for the direct decay of the 125 GeV Higgs boson to fermions. **NATURE PHYSICS** 10: pp. 557-560. (2014)
2. Bodwin GT et al. incl. [Bencze Gy](#), [Hajdu Cs](#), [Hidas P](#), [Horvath D](#), [Sikler F](#), [Veszpremi V](#), [Vesztergombi Gy](#), [Zsigmond AJ](#) [2200 authors]: Fragmentation contributions to  $J/\psi$  production at the Tevatron and the LHC. **PHYS REV LETT**, 113:(2) Paper 022001. (2014)
3. Chatrchyan S et al. incl. [Bencze Gy](#), [Hajdu Cs](#), [Hidas P](#), [Horvath D](#), [Sikler F](#), [Veszpremi V](#), [Vesztergombi Gy](#), [Zsigmond AJ](#) [2200 authors]: Search for top squark and higgsino production using diphoton higgs boson decays. **PHYS REV LETT**, 112:(16) Paper 161802. 15 p. (2014)
4. Chatrchyan S et al. incl. [Bencze Gy](#), [Hajdu Cs](#), [Hidas P](#), [Horvath D](#), [Sikler F](#), [Veszpremi V](#), [Vesztergombi Gy](#), [Zsigmond AJ](#) [2202 authors]: Search for top-quark partners with charge 5/3 in the same-sign dilepton final state. **PHYS REV LETT**, 112:(17) Paper 171801. 16 p. (2014)
5. Chatrchyan S et al. incl. [Bencze Gy](#), [Hajdu Cs](#), [Hidas P](#), [Horvath D](#), [Sikler F](#), [Veszpremi V](#), [Vesztergombi Gy](#), [Zsigmond AJ](#) [2200 authors]: Search for flavor-changing neutral currents in top-quark decays  $t \rightarrow zq$  in pp collisions at  $\sqrt{s} = 8\text{TeV}$ . **PHYS REV LETT**, 112:(17) Paper 171802. 16 p. (2014)
6. Chatrchyan S et al. incl. [Bencze Gy](#), [Hajdu Cs](#), [Hidas P](#), [Horvath D](#), [Sikler F](#), [Veszpremi V](#), [Vesztergombi Gy](#), [Zsigmond AJ](#) [2202 authors]: Measurements of  $t \bar{t}$  spin correlations and top-quark polarization using dilepton final states in pp collisions at  $\sqrt{s} = 7\text{TeV}$ . **PHYS REV LETT**, 112:(18) Paper 182001. 16 p. (2014)
7. Chatrchyan S et al. incl. [Bencze Gy](#), [Hajdu Cs](#), [Hidas P](#), [Horvath D](#), [Sikler F](#), [Veszpremi V](#), [Vesztergombi Gy](#), [Zsigmond AJ](#) [2209 authors]: Measurement of inclusive W and Z boson production cross sections in pp collisions at  $\sqrt{s} = 8\text{TeV}$ . **PHYS REV LETT**, 112:(19) Paper 191802. 17 p. (2014)
8. Chatrchyan S et al. incl. [Bencze Gy](#), [Hajdu Cs](#), [Hidas P](#), [Horvath D](#), [Sikler F](#), [Veszpremi V](#), [Vesztergombi Gy](#), [Zsigmond AJ](#) [2207 authors]: Observation of the associated

production of a single top quark -and a W boson in  $pp$  collisions at  $\sqrt{s} = 8\text{TeV}$ . **PHYS REV LETT**, 112:(23) Paper 231802. 16 p. (2014)

9. Chatrchyan S et al. incl. [Bencze Gy](#), [Hajdu Cs](#), [Hidas P](#), [Horvath D](#), [Sikler F](#), [Veszpremi V](#), [Vesztergombi Gy](#), [Zsigmond AJ](#) [2206 authors]: Evidence of  $b$ -Jet Quenching in PbPb Collisions at  $\sqrt{s_{NN}} = 2.76\text{ TeV}$ . **PHYS REV LETT**, 113: Paper 132301. 16 p. (2014)
10. Khachatryan V et al. incl. [Bencze Gy](#), [Hajdu Cs](#), [Hidas P](#), [Horvath D](#), [Sikler F](#), [Veszpremi V](#), [Vesztergombi Gy](#), [Zsigmond AJ](#) [2147 authors]: Measurement of Prompt  $\psi(2S) \rightarrow J/\psi$  Yield Ratios in Pb-Pb and p-p Collisions at  $\sqrt{s_{NN}} = 2.76\text{ TeV}$ . **PHYS REV LETT**, 113:(26) Paper 262301. 15 p. (2014)
11. Chatrchyan S et al. incl. [Bencze Gy](#), [Hajdu Cs](#), [Hidas P](#), [Horvath D](#), [Sikler F](#), [Veszpremi V](#), [Vesztergombi Gy](#), [Zsigmond AJ](#) [2231 authors]: Evidence for the 125 GeV Higgs boson decaying to a pair of  $\tau$  leptons. **J HIGH ENERGY PHYS**, 2014:(5) Paper 104. 72 p. (2014)
12. Chatrchyan S et al. incl. [Bencze Gy](#), [Hajdu Cs](#), [Hidas P](#), [Horvath D](#), [Sikler F](#), [Veszpremi V](#), [Vesztergombi Gy](#), [Zsigmond AJ](#) [2207 authors]: Measurement of Higgs boson production and properties in the WW decay channel with leptonic final states. **J HIGH ENERGY PHYS**, 2014:(1) Paper 096. 86 p. (2014)
13. Chatrchyan S et al. incl. [Bencze Gy](#), [Hajdu Cs](#), [Hidas P](#), [Horvath D](#), [Sikler F](#), [Veszpremi V](#), [Vesztergombi Gy](#), [Zsigmond AJ](#) [2210 authors]: Search for new physics in events with same-sign dileptons and jets in pp collisions at  $\sqrt{s} = 8\text{ TeV}$ . **J HIGH ENERGY PHYS**, 2014:(1) Paper 163. (2014)
14. Chatrchyan S et al. incl. [Bencze Gy](#), [Hajdu Cs](#), [Hidas P](#), [Horvath D](#), [Sikler F](#), [Veszpremi V](#), [Vesztergombi Gy](#), [Zsigmond AJ](#) [2202 authors]: Studies of azimuthal dihadron correlations in ultra-central PbPb collisions at  $\sqrt{s_{NN}} = 2.76\text{ TeV}$ . **J HIGH ENERGY PHYS**, 2014:(2) Paper 088. 38 p. (2014)
15. Chatrchyan S et al. incl. [Bencze Gy](#), [Hajdu Cs](#), [Hidas P](#), [Horvath D](#), [Sikler F](#), [Veszpremi V](#), [Vesztergombi Gy](#), [Zsigmond AJ](#) [2210 authors]: Measurement of the  $t\bar{t}$  production cross section in the dilepton channel in pp collisions at  $\sqrt{s} = 8\text{ TeV}$ . **J HIGH ENERGY PHYS**, 2014:(2) Paper 024. 31 p. (2014)
16. Chatrchyan S et al. incl. [Bencze Gy](#), [Hajdu Cs](#), [Hidas P](#), [Horvath D](#), [Sikler F](#), [Veszpremi V](#), [Vesztergombi Gy](#), [Zsigmond AJ](#) [2207 authors]: Study of double parton scattering using W + 2-jet events in proton-proton collisions at  $\sqrt{s} = 7\text{ TeV}$ . **J HIGH ENERGY PHYS**, 2014:(3) Paper 032. 45 p. (2014)
17. Chatrchyan S et al. incl. [Bencze Gy](#), [Hajdu Cs](#), [Hidas P](#), [Horvath D](#), [Sikler F](#), [Veszpremi V](#), [Vesztergombi Gy](#), [Zsigmond AJ](#) [2207 authors]: Search for  $W' \rightarrow t\bar{b}$  decays in the lepton plus jets final state in pp collisions at  $\sqrt{s} = 8\text{ TeV}$ . **J HIGH ENERGY PHYS**, 2014:(5) Paper 108. 35 p. (2014)
18. Chatrchyan S et al. incl. [Bencze Gy](#), [Hajdu Cs](#), [Hidas P](#), [Horvath D](#), [Sikler F](#), [Veszpremi V](#), [Vesztergombi Gy](#), [Zsigmond AJ](#) [2224 authors]: Measurements of the  $t\bar{t}$  charge asymmetry using the dilepton decay channel in pp collisions at  $\sqrt{s} = 7\text{ TeV}$ . **J HIGH**

**ENERGY PHYS**, 2014:(4) Paper 191. 35 p. (2014)

19. Chatrchyan S et al. incl. [Bencze Gy](#), [Hajdu Cs](#), [Hidas P](#), [Horvath D](#), [Sikler F](#), [Veszpremi V](#), [Vesztergombi Gy](#), [Zsigmond AJ](#) [2209 authors]: Event activity dependence of  $Y$  ( $nS$ ) production in  $\sqrt{s_{NN}} = 5.02$  TeV pPb and  $\sqrt{s} = 2.76$  TeV pp collisions. **J HIGH ENERGY PHYS**, 2014:(4) Paper 103. 37 p. (2014)
20. Chatrchyan S et al. incl. [Bencze Gy](#), [Hajdu Cs](#), [Hidas P](#), [Horvath D](#), [Sikler F](#), [Veszpremi V](#), [Vesztergombi Gy](#), [Zsigmond AJ](#) [2224 authors]: Measurement of associated  $W$  plus charm production in pp collisions at  $\sqrt{s} = 7$  TeV. **J HIGH ENERGY PHYS**, 2014:(2) Paper 013. 63 p. (2014)
21. Chatrchyan S et al. incl. [Bencze Gy](#), [Hajdu Cs](#), [Hidas P](#), [Horvath D](#), [Sikler F](#), [Veszpremi V](#), [Vesztergombi Gy](#), [Zsigmond AJ](#) [2221 authors]: Measurement of the production cross sections for a  $Z$  boson and one or more  $b$  jets in pp collisions at  $\sqrt{s} = 7$  TeV. **J HIGH ENERGY PHYS**, 2014:(6) Paper 120. 39 p. (2014)
22. Chatrchyan S et al. incl. [Bencze Gy](#), [Hajdu Cs](#), [Hidas P](#), [Horvath D](#), [Sikler F](#), [Veszpremi V](#), [Vesztergombi Gy](#), [Zsigmond AJ](#) [2221 authors]: Search for pair production of excited top quarks in the lepton + jets final state. **J HIGH ENERGY PHYS**, 2014:(6) Paper 125. 35 p. (2014)
23. Chatrchyan S et al. incl. [Bencze Gy](#), [Hajdu Cs](#), [Hidas P](#), [Horvath D](#), [Sikler F](#), [Veszpremi V](#), [Vesztergombi Gy](#), [Zsigmond AJ](#) [2214 authors]: Search for new physics in the multijet and missing transverse momentum final state in proton-proton collisions at  $\sqrt{s} = 8$  TeV. **J HIGH ENERGY PHYS**, 2014:(6) Paper 055. 38 p. (2014)
24. Chatrchyan S et al. incl. [Bencze Gy](#), [Hajdu Cs](#), [Hidas P](#), [Horvath D](#), [Sikler F](#), [Veszpremi V](#), [Vesztergombi Gy](#), [Zsigmond AJ](#) [2223 authors]: Measurement of the triple-differential cross section for photon + jets production in proton-proton collisions at  $\sqrt{s} = 7$  TeV. **J HIGH ENERGY PHYS**, 2014:(6) Paper 009. 35 p. (2014)
25. Khachatryan V et al. incl. [Bencze Gy](#), [Hajdu Cs](#), [Hidas P](#), [Horvath D](#), [Sikler F](#), [Veszpremi V](#), [Vesztergombi Gy](#), [Zsigmond AJ](#) [2126 authors]: Measurement of the  $t$ -channel single-top-quark production cross section and of the  $|V_{tb}|$  CKM matrix element in pp collisions at  $\sqrt{s} = 8$  TeV. **J HIGH ENERGY PHYS**, 2014:(6) Paper 090. 43 p. (2014)
26. Khachatryan V et al. incl. [Bencze Gy](#), [Hajdu Cs](#), [Hidas P](#), [Horvath D](#), [Sikler F](#), [Veszpremi V](#), [Vesztergombi Gy](#), [Zsigmond AJ](#) [2133 authors]: Search for massive resonances decaying into pairs of boosted bosons in semi-leptonic final states at  $\sqrt{s} = 8$  TeV. **J HIGH ENERGY PHYS**, 2014:(8) Paper 174. 59 p. (2014)
27. Khachatryan V et al. incl. [Bencze Gy](#), [Hajdu Cs](#), [Hidas P](#), [Horvath D](#), [Sikler F](#), [Veszpremi V](#), [Vesztergombi Gy](#), [Zsigmond AJ](#) [2129 authors]: Measurement of prompt  $J/\psi$  pair production in pp collisions at  $\sqrt{s} = 7$  TeV. **J HIGH ENERGY PHYS**, 2014:(9) Paper 094. 35 p. (2014)
28. Khachatryan V et al. incl. [Bencze Gy](#), [Hajdu Cs](#), [Hidas P](#), [Horvath D](#), [Sikler F](#), [Veszpremi V](#), [Vesztergombi Gy](#), [Zsigmond AJ](#) [2137 authors]: Search for the associated production of the Higgs boson with a top-quark pair. **J HIGH ENERGY PHYS**,

2014:(9) Paper 87. 64 p. (2014)

29. Khachatryan V et al. incl. Bencze Gy, Hajdu Cs, Hidas P, Horvath D, Sikler F, Veszpremi V, Vesztergombi Gy, Zsigmond AJ [2132 authors]: Study of hadronic event-shape variables in multijet final states in pp collisions at  $\sqrt{s} = 7$  TeV. **J HIGH ENERGY PHYS**, 2014:(10) Paper 087. 34 p. (2014)
30. Khachatryan V et al. incl. Bencze Gy, Hajdu Cs, Hidas P, Horvath D, Sikler F, Veszpremi V, Vesztergombi Gy, Zsigmond AJ [2147 authors]: Identification techniques for highly boosted W bosons that decay into hadrons. **J HIGH ENERGY PHYS**, 2014:(12) Paper 17. 46 p. (2014)

### **NA49 Collaboration**

1. Anticic T et al. incl. Barna D, Fodor Z, Laszlo A, Palla G, Sikler F, Vesztergombi Gy [68 authors]: Phase-space dependence of particle-ratio fluctuations in Pb+Pb collisions from 20A to 158A GeV beam energy. **PHYS REV C**, 89:(5) Paper 054902. 28 p. (2014)

### **FOPI Collaboration**

1. Zinyuk V et al. incl. Fodor Z, Kecskeméti KJ, Seres Z [65 authors]: Azimuthal emission patterns of K<sup>+</sup> and of K<sup>-</sup> mesons in Ni plus Ni collisions near the strangeness production threshold. **PHYS REV C**, 90:(2) Paper 025210. 6 p. (2014)

***See also: R-B. ALICE Collaboration, R-I. NA61/SHINE Collaboration***